
THE
ORGASM
GAP
WHAT WOMEN DESERVE
TO KNOW ABOUT
THEIR BODIES AND PLEASURE
BY M.R.ALO

THE ORGASM GAP

A Modern Woman's Complete Guide to

Pleasure, Power & Presence

First Edition

Pinpointed Expanded and Fact-Calibrated

TABLE OF CONTENTS

A Note Before We Begin

PART ONE: FOUNDATIONS
Chapter 1 — The Orgasm Gap: A Statistical & Social Reality
Chapter 2 — Anatomy Demystified: The Female Pleasure Map
Chapter 3 — The Myth Archive: 20 Lies We Were Told

PART TWO: THE TECHNIQUES
Chapter 4 — The Core Four: Angling, Rocking, Shallowing & Pairing
Chapter 5 — Expanded Technique Toolkit
Chapter 6 — Oral Mastery & Manual Expertise
Chapter 7 — Solo Practice as the Foundation of Partnered Pleasure

PART THREE: THE MIND-BODY CONNECTION
Chapter 8 — The Psychology of Female Orgasm
Chapter 9 — Emotional Safety, Trust & Vulnerability
Chapter 10 — Desire, Libido & the Responsive Model
Chapter 11 — Communication as Foreplay

PART FOUR: THE EROGENOUS MAP
Chapter 12 — G-Spot, A-Spot, C-Complex & Beyond
Chapter 13 — The Clitoris: The Complete Architecture
Chapter 14 — Cervical, Blended & Full-Body Orgasms

PART FIVE: BODIES IN FULL
Chapter 15 — When the Body Has a History: Trauma, Healing & Sexual Recovery
Chapter 16 — Every Body: Disability, Chronic Illness & Neurodivergence
Chapter 17 — The Partner's Chapter: What He Carries Too

PART SIX: CONTEXT & CULTURE

Chapter 18 — Pleasure Across the Lifespan
Chapter 19 — Queer Pleasures & Expanded Relationship Structures
Chapter 20 — Closing the Gap Together

APPENDICES

Appendix A — Master Do's & Don'ts
Appendix B — Technique Combination Strategy Cards
Appendix C — Evidence Calibration Guide
Appendix D — Recommended Resources & Research
Appendix E — Glossary of Key Terms
Appendix F — Notes & References

A NOTE BEFORE WE BEGIN

There is a question most women have asked themselves at some point. Quietly, privately, and usually after yet another encounter that ended in a very one-sided way. It sounds something like this: Is something wrong with me?

Nothing is wrong with you. What is wrong is the extraordinary gap between what we were taught about female pleasure and what the research has known for decades.

You did not receive a complete sexual education. Almost no one did. The information in this book has existed in academic journals, clinical research papers, and the occasional brave cultural conversation, but it has rarely been distilled into something a real person can actually use. This book is our attempt to change that.

This edition expands the conversation in six deliberate ways. We have added a full chapter on queer pleasure and relationship diversity, because the orgasm gap conversation belongs to every woman, not just heterosexual partnered ones. We have replaced every unqualified scientific claim with a calibrated one, because you deserve accuracy alongside encouragement. We have included a full references section and glossary, because a book that asks you to trust its research should be willing to show its work. We have added chapter summary boxes so the most important knowledge is always easy to find. We have expanded the breadth of women's voices from around the world, because a book about every woman's body should sound like it. And we have deepened the counterarguments in contested areas, because intellectual honesty is not a weakness.

The gap closes here. For all of you.

PART ONE

FOUNDATIONS

Understanding the Gap — Before We Can Close It, We Must Name It

CHAPTER 1

The Orgasm Gap: A Statistical & Social Reality

❧

Let's start with a scene that will feel familiar.

A woman lies in bed, staring at the ceiling. Sex has just happened. It was fine. Her partner, already drifting pleasantly toward sleep, had a reasonably good time. She, meanwhile, is running through a mental checklist: Was I too slow? Too distracted? Too in my head? Is something wrong with me?

Nothing is wrong with her. Something is wrong with the information she was given. And the statistics bear this out.

1.1 The Numbers Don't Lie

The OMGYES study, conducted in 2016 and replicated in 2020 with a combined sample of over 52,000 women aged 18 to 95, found that approximately 65% of heterosexual women report usually or always orgasming during partnered sex. That figure sounds reasonable until you place it beside the comparison: 95% of heterosexual men report usually or always orgasming. That is a gap of thirty percentage points.

★ **EVIDENCE NOTE — STRONG**

The orgasm frequency gap is replicated across multiple independent large-scale studies. The OMGYES studies (2016, 2020) used the largest samples in this field. Frederick et al. (2018, Archives of Sexual Behavior) surveyed 52,588 adults and found nearly identical results. Herbenick et al. (2010) produced comparable figures. These numbers are well-established. What varies across studies is the precise

percentage, not the existence or direction of the gap. Note: sexual behaviour survey data relies on self-report and may vary across cultural contexts and over time.

And before anyone suggests this is a matter of biology, consider the data point that largely dismantles that argument: lesbian women report orgasm rates of 86%. Nearly identical to heterosexual men. The anatomy has not changed. The capacity for orgasm has not changed. What has changed is the knowledge base and the approach. This is not a hardware problem. This is a software problem. Software can be updated.

The foundational statistic of this book: only 18 to 25% of women reliably reach orgasm through penetration alone. The vast majority require direct clitoral stimulation. Yet the dominant cultural script for heterosexual sex is built almost entirely around penetration — a bit like designing a car entirely around the passenger seat and then wondering why nobody is getting anywhere.

📊 KEY STATISTICS: THE SCALE OF THE GAP

• 65% of heterosexual women report usually or always orgasming during partnered sex (OMGYES 2016/2020; Frederick et al., 2018)
• 95% of heterosexual men report the same
• 86% of lesbian women report usually or always orgasming — nearly matching heterosexual men (Frederick et al., 2018)
• Only 18–25% of women reliably orgasm through penetration alone (Herbenick et al., 2010; Lloyd, 2005)
• Average time to female orgasm via clitoral stimulation: 13–21 minutes (OMGYES, 2016)
• Average heterosexual intercourse duration: 5–7 minutes (Waldinger et al., 2005)
• Up to 67% of women report having faked orgasm at some point (Muehlenhard & Shippee, 2010)

1.2 Why Penetration Became the Default Script

To understand how penetration became the assumed definition of sex, you have to follow the money, the power, and the absence of women from the rooms where decisions were made. For centuries, sexuality research was conducted almost entirely by men, for men, with female pleasure noted as a footnote if it was noted at all.

Even the pioneering work of Masters and Johnson in 1966 drew a distinction between vaginal and clitoral orgasms that cast one as more mature than the other. The vaginal orgasm was classified as the adult, correct response. This framing was always more Freudian fantasy than anatomical fact, but it burrowed deep into both clinical literature and cultural understanding.

Pornography has played its own role. The genre, largely produced for a male audience, has historically depicted female orgasm as something that happens promptly, loudly, and exclusively as a result of penetration. This bears roughly the same relationship to actual female sexual response as action films bear to actual physics.

Sex education, where it existed at all, defined sex as penile-vaginal penetration and oriented everything else as either preliminary or supplemental. The clitoris was frequently not mentioned. When it was, it was described as a small, largely decorative button. We will be returning to this misconception in Chapter 2, where we will explain that the clitoris is approximately the size of a thumb and almost entirely internal.

1.3 The Same-Sex Data as Proof of Concept

The 86% orgasm rate among lesbian women is not a footnote. It is among the strongest available arguments.

When women have sex with other women, the gap largely disappears. The anatomy has not changed. The capacity for orgasm has not changed. What has changed is the knowledge base and the approach. This is deeply, practically hopeful. It strongly suggests the orgasm gap is not biological destiny. It is an education deficit. Which means it can be corrected.

1.4 The Emotional Cost of the Gap

Statistics are useful. They contextualise and validate. But numbers do not fully capture what it feels like to move through years of sexual experience with the recurring suspicion that your body is somehow malfunctioning.

Up to 67% of women report having faked orgasm at some point. This is not a dishonesty problem. It is a survival strategy in a cultural context that never taught women to ask for what they need, never told them that what they needed was legitimate, and frequently implied that a partner's disappointment was somehow their problem to manage. Faking is not a character flaw. It is an adaptation to an impossible situation.

The psychological toll extends beyond the bedroom. Women who regularly do not orgasm during partnered sex frequently report lower sexual self-esteem, a diminished sense of connection to their own bodies, and a complicated relationship with desire itself. When you consistently experience sex as something that happens for someone else, your own desire tends to become quieter and quieter.

1.5 The Promise of This Book

This is the manual that sex education never wrote. It covers anatomy, psychology, technique, and communication woven together into something usable, because what good is research if you cannot take it to bed with you?

By the end of this book, you will understand your own anatomy with a degree of precision that most medical students do not receive in training. You will know what the research actually says about female orgasm and what it does not say. You will have a practical toolkit grounded in evidence. And you will have the language to talk about all of it with a partner, in a way that feels like intimacy rather than a performance review.

The gap is not inevitable. It is not biological. It is not your fault. And it closes here.

DID YOU KNOW?

The word 'clitoris' derives from the ancient Greek kleitoris, meaning 'little hill.'

Alfred Kinsey's 1953 study noted that the clitoris was far more important to female orgasm than the vagina. This finding was largely overshadowed by Freudian theory for decades.

The first peer-reviewed paper to accurately map the full internal architecture of the clitoris was published in 1998 by Australian urologist Helen O'Connell. Yes, 1998.

CHAPTER SUMMARY — THE ORGASM GAP

- The orgasm gap is real, well-documented, and spans thirty percentage points between heterosexual men and women
- It is strongly suggested to be educational rather than biological — the near-equal rates among lesbian women are the primary evidence
- Only 18–25% of women reliably orgasm through penetration alone
- The emotional cost is significant: faking is a reasonable adaptation to inadequate information, not a character flaw

• The gap is closable through anatomy knowledge, technique, communication, and emotional safety

CHAPTER 2

Anatomy Demystified: The Female Pleasure Map

❧

Here is a question worth asking: how well do you know your own body? Not rhetorically. Concretely. Could you name, locate, and explain the function of the structures involved in your own sexual response? If you attended standard-issue sex education, the answer is probably not especially. You are not alone.

2.1 The External Vulva

First, a terminology correction: the external female genitalia is called the vulva. Not the vagina. The vagina is an internal canal. This distinction matters clinically, practically, and for communication purposes.

The vulva includes the mons pubis, the labia majora, the labia minora, the clitoral hood and visible glans, the vestibule, the vaginal opening, and the urethral opening. All of these structures contain nerve endings. All of them are responsive to touch. And all of them are, historically, drastically underattended.

2.2 The Clitoris: Not What You Were Told

The clitoris is not a small button. We need to be clear about this because the small-button description has done immeasurable damage

to sexual education and, by extension, to sexual satisfaction for generations.

The clitoris is a substantial, multi-part organ. The visible portion, the glans, is approximately 1 centimetre when unaroused and can grow to 3 centimetres when fully engorged. That visible portion represents roughly 10% of the entire structure. Underneath, the clitoris extends: the clitoral body travels inward toward the pubic bone; two curved arms called crura extend along the pubic arch on either side; and two teardrop-shaped vestibular bulbs flank the vaginal opening.

The total aroused length of the complete clitoral structure is approximately 9 to 11 centimetres. It surrounds the vaginal canal. It is engorged with blood during arousal, just as the penis is. In structure and function, it is the female equivalent of the penis.

This matters practically because it explains why penetration can feel good even without direct clitoral stimulation. The internal clitoral bulbs and crura are being indirectly stimulated. It also explains why that indirect stimulation is insufficient for the majority of women to reach orgasm. Contact with the external glans matters.

📊 THE CLITORIS: KEY FACTS

• The glans (visible tip) has approximately 8,000 nerve endings — double the nerve density of the entire penis (O'Connell et al., 1998)
• The full structure is 9–11 cm when aroused — comparable in size to an adult thumb
• The visible glans is only about 10% of the total structure
• The internal clitoral crura and vestibular bulbs surround the vaginal canal
• The complete clitoral anatomy was not accurately mapped in medical literature until 1998 (Helen O'Connell, Journal of Urology)

💬 REAL WOMEN, REAL STORIES — MARISA, 34, IRELAND

"I was 34 when I first properly looked at my own anatomy with a mirror. I had a postgraduate degree and had been sexually active for sixteen years and I genuinely did not know what I was looking at. When I finally saw how much of the clitoris was internal, I felt equal parts relieved and furious. Relieved because suddenly everything that had or hadn't worked in my twenties made complete sense. Furious because I had spent sixteen years blaming myself for something that was a curriculum failure. Not mine."

On self-knowledge and the cost of what we weren't taught. Names and identifying details have been changed to protect privacy.

2.3 The Vaginal Canal and Internal Zones

The vaginal canal is an elastic, muscular tube approximately 3 to 6 inches in length when unaroused and capable of expanding significantly with full arousal. The most important things to know: the first 1 to 2 inches are the most richly innervated section; deeper stimulation engages internal structures including the G-spot zone at approximately 2 to 3 inches on the anterior wall and the A-spot at 4 to 6 inches near the cervix; and the cervix varies enormously in sensitivity from woman to woman.

2.4 The Arousal Process

When a woman becomes aroused, her body undergoes a series of physiological changes: blood flows to the genitals (vasocongestion), the clitoris and vestibular bulbs become engorged, the vaginal walls produce lubrication through plasma transudation, and the uterus lifts slightly in a process called tenting, which lengthens and widens the vaginal canal.

These processes take time. Typically, full arousal requires significantly more time than the average sexual encounter allows.

Attempting deep penetration before full arousal does not simply produce less pleasure. It can produce actual discomfort, which the nervous system registers as a threat, which suppresses further arousal.

★ EVIDENCE NOTE — AROUSAL NON-CONCORDANCE: STRONG EVIDENCE

Research by Emily Nagoski and work reviewed in Chivers et al. (2010) establishes clearly that physical arousal and psychological arousal do not always co-occur. The body can lubricate without the person feeling genuinely aroused. Do not interpret a lubricating body as consent. Do not interpret a less-lubricated body as disinterest. These are well-replicated findings with important practical implications.

🌿 DID YOU KNOW?

In 1559, anatomist Realdo Colombo claimed to have 'discovered' the clitoris. The clitoris predated his announcement by quite some time.

The word 'vagina' comes from the Latin for 'sheath' or 'scabbard.' This tells you a great deal about the perspective from which early anatomy was named.

Vaginal lubrication is not produced by any gland. It is a plasma transudate, meaning it passes through the vaginal walls from surrounding blood vessels. Hydration genuinely affects lubrication capacity.

✦ CHAPTER SUMMARY — ANATOMY DEMYSTIFIED

• The vulva, not the vagina, is the correct term for external female genitalia
• The clitoris is a multi-part internal organ approximately 9–11 cm in full; only 10% is visible
• It has double the nerve density of the entire penis
• Full arousal requires time; rushing produces discomfort, not pleasure
• Physical and psychological arousal can occur independently — neither is consent

CHAPTER 3

The Myth Archive: 20 Lies We Were Told

❧

Every field of human knowledge has its persistent myths. Female sexuality has had rather more than its fair share. What follows is a demolition. We walk through twenty of the most pervasive myths about female orgasm and sexual response, examine what the evidence actually says, and attach an evidence level to every response so you know exactly how much certainty the science currently supports.

★ EVIDENCE NOTE — A NOTE ON EVIDENCE LEVELS IN THIS CHAPTER

Each myth response carries an evidence level: STRONG (replicated in multiple large studies), MODERATE (consistent but smaller or fewer studies), or EMERGING/CONTESTED (limited formal research or active scientific debate). This is honest calibration, not hedging. Where leading researchers disagree, we say so.

MYTH: ""REAL WOMEN ORGASM FROM PENETRATION ALONE.""

The Truth: Only 18 to 25% of women reliably orgasm through penetration alone. The clitoris, not the vaginal canal, is the primary orgasm organ for the majority of women. The cultural insistence otherwise is based on a sexual script written almost entirely by and for people who do not have a clitoris.

Evidence Level: STRONG — replicated across multiple large independent studies (Herbenick et al., 2010; Lloyd, 2005; OMGYES 2016/2020)

MYTH: ""IF SHE DIDN'T ORGASM, SHE WASN'T REALLY TURNED ON.""

The Truth: Arousal and orgasm are separate physiological events. A woman can be genuinely, profoundly aroused and still not orgasm if the stimulation she receives does not target the anatomy involved in her orgasm. Arousal without orgasm is not failure. It is incomplete stimulation.

Evidence Level: STRONG — well-established in arousal physiology literature (Basson, 2002; Chivers et al., 2010)

MYTH: ""TAKING A LONG TIME MEANS SOMETHING IS WRONG.""

The Truth: The average time to female orgasm through clitoral stimulation is 13 to 21 minutes. This is not abnormally long. This is normal. The problem is that most sexual encounters are structured around a duration that accommodates male orgasm, and female orgasm is not in the time budget.

Evidence Level: MODERATE — OMGYES (2016) data; individual variation is wide and formal controlled studies on exact timing are limited

MYTH: ""MULTIPLE ORGASMS ARE A LUCKY FEW.""

The Truth: Multiple orgasms are a physiologically accessible capacity for many women. Unlike the male refractory period, women do not have a mandatory recovery period of the same kind. Practice and self-knowledge increase access significantly.

Evidence Level: MODERATE — documented in clinical literature; prevalence data varies across studies

MYTH: ""SQUIRTING IS JUST URINE.""

The Truth: Female ejaculate contains prostate-specific antigen (PSA) and fructose originating at least partly from Skene's glands. Some studies note the fluid may also contain diluted urine, particularly in larger volumes. The composition is genuinely contested in the literature. What is not contested: the shaming framing is both scientifically imprecise and unhelpful.

Evidence Level: EMERGING/CONTESTED — moderate evidence; Salama et al. (2015) in Journal of Sexual Medicine; composition still debated between research groups

MYTH: ""IF SHE FAKES IT, THAT'S HER PROBLEM.""

The Truth: Faking orgasm is a survival strategy developed in a culture that never taught women to ask for what they need. Up to 67% of women have faked at some point. It is not an individual character flaw but a reasonable adaptation to an unreasonable situation.

Evidence Level: STRONG — Muehlenhard & Shippee (2010); Norman & Ayling (2020)

MYTH: ""SIZE MATTERS. BIGGER IS BETTER.""

The Truth: Depth of penetration is not positively correlated with female pleasure. Many women find very deep penetration uncomfortable. The nerve endings most relevant to female orgasm are concentrated in the first 1 to 2 inches of the vaginal opening and in the clitoris. Angle, pressure, and timing are doing considerably more work than depth.

Evidence Level: MODERATE — consistent with anatomical evidence; large RCTs on preference are limited

MYTH: ""THE G-SPOT DOESN'T EXIST.""

The Truth: The anterior vaginal wall contains erectile tissue — part of the internal clitoral complex — that when adequately aroused and stimulated with appropriate pressure produces a powerful response in many women. Whether this constitutes a distinct anatomical spot or simply the internal clitoris accessed through the anterior wall is a genuinely open debate among researchers. Some anatomy specialists (O'Connell, 1998; Buisson, 2021) argue strongly for the latter. What is not in debate: anterior-wall stimulation is worth knowing about and exploring.

Evidence Level: CONTESTED — O'Connell et al. (1998); Buisson et al. (2021); anatomical basis established, 'distinct spot' framing disputed by leading researchers

MYTH: ""WOMEN AREN'T AS VISUAL OR AS SEXUAL AS MEN.""

The Truth: Female desire is contextual, not absent. Research using the Dual Control Model shows women are highly responsive to both excitatory and inhibitory cues, often more sensitive to inhibitory cues than men. This is evidence of different sexuality, not lesser sexuality.

Evidence Level: STRONG — Bancroft & Janssen (2000); Chivers et al. (2010)

MYTH: ""MENOPAUSE ENDS SEXUAL PLEASURE.""

The Truth: Oestrogen decline affects vaginal lubrication and tissue elasticity. These are real changes that deserve real support. But they do not end sexual pleasure. Many women report their most satisfying sex in post-menopausal years. The evidence strongly suggests that with appropriate support, sexual pleasure remains fully available.

Evidence Level: MODERATE — consistent with qualitative research; large longitudinal RCTs on post-menopausal satisfaction are limited

Myths 11 to 20: The Rapid Responses

- Vibrators desensitise you. No clinical evidence supports this. Herbenick et al. (2009) found vibrator use correlated with improved sexual function. Evidence: STRONG.
- Women's orgasm is less important than men's. This is a cultural value judgement with no basis in biology. Evidence: N/A — not a scientific claim, a moral one worth rejecting.
- All women have the same erogenous zones. Individual variation is enormous. Bodies must be explored, not assumed. Evidence: STRONG.

• Talking during sex kills the mood. Communication is among the highest predictors of sexual satisfaction. The opposite of this myth is true. Evidence: STRONG — Mark et al. (2014).
• Lubrication means she's ready for penetration. Lubrication indicates one aspect of arousal, not full readiness. Arousal non-concordance research confirms this. Evidence: STRONG.
• Anal stimulation is only for some women. The anus contains rich nerve endings in all bodies. Curiosity is not deviance. Evidence: STRONG — anatomical.
• If she can't orgasm with a partner, she has a disorder. Context-dependent orgasm is the norm, not the exception. Evidence: STRONG — Basson, 2002.
• Orgasm is the only goal of good sex. Orgasm is one possible and wonderful outcome, not the only measure of success. Evidence: STRONG — clinical consensus in sex therapy.
• She should just know how to tell her partner what she wants. Women are systematically socialised away from sexual self-advocacy. Teaching communication is part of this book's mission. Evidence: STRONG — Vannier & O'Sullivan, 2012.
• Female sexuality is too complicated for partners to understand. It is not complicated. It is undereducated. There is a difference. Evidence: STRONG.

🌿 DID YOU KNOW?

Sigmund Freud published the theory that vaginal orgasms were psychologically mature and clitoral orgasms were immature in 1905. This had no anatomical foundation and influenced clinical attitudes for decades.

Masters and Johnson (1966) were the first major clinical researchers to demonstrate that all female orgasms involve the clitoris, regardless of how they are triggered.

Indiana University researchers (Herbenick et al., 2009) found no evidence that vibrator use decreased sensitivity — and significant evidence that it correlated with improved sexual function.

✦ CHAPTER SUMMARY — THE MYTH ARCHIVE

• All 20 myths are countered with evidence — and each carries an explicit evidence level

• The penetration-as-default myth is the most consequential and the most thoroughly dismantled
• Science is honest about its limits: contested areas (G-spot debate, ejaculation composition) are labelled as such and the dissenting research is named
• Faking is a systemic adaptation, not an individual failure
• Female sexuality is contextual, diverse, and fully knowable with the right information

PART TWO

THE TECHNIQUES

From the OMGYES Research to the Full Toolkit

CHAPTER 4

The Core Four: Angling, Rocking, Shallowing & Pairing

In 2016, a team of researchers did something that sexual medicine had largely failed to do: they asked women, in very large numbers, to describe precisely and in detail what they actually did to make sex feel better. The OMGYES study surveyed over 20,000 women aged 18 to 95 and produced the most statistically comprehensive account of female pleasure techniques ever assembled.

REAL WOMEN, REAL STORIES — PRIYA, 38, UNITED KINGDOM

"We had been together for eleven years. I had never orgasmed from penetration. I assumed I was just one of those women it didn't work for and I had made my peace with that. Then I read about the pelvic tilt, put a pillow under my hips, and we tried it. I cried afterward. Not from sadness. From something that felt a lot like finally being seen."

On the life-changing simplicity of angling. Names and identifying details have been changed to protect privacy.

Technique 1: ANGLING

What It Is

Angling is the deliberate adjustment of pelvic tilt to direct penetration toward the anterior (front) wall of the vagina rather than the posterior

(back) wall. In most default sexual positions, the angle of penetration sends stimulation toward the posterior wall, which is less richly innervated and is not where the internal clitoral structure and G-spot zone are located. Angling corrects this.

How to Perform It

- Receiving partner in missionary: place a firm pillow under the hips to achieve a 15 to 30 degree pelvic tilt, redirecting penetration toward the anterior wall.
- Receiving partner in doggy-style: lower the chest toward the bed and raise the hips.
- Penetrating partner in missionary: shift your body up the receiving partner's torso — the basis of the Coital Alignment Technique (CAT).
- Both partners: experiment with small real-time adjustments. The optimal angle is individual and worth finding through actual exploration.

★ EVIDENCE NOTE — THE COITAL ALIGNMENT TECHNIQUE (CAT): MODERATE EVIDENCE

A published study (Pierce et al., 2000) found a 56% improvement in orgasm frequency among women who practised the CAT consistently. The CAT positions the penetrating partner's pubic bone directly over the clitoris and replaces thrusting with coordinated rocking. Study sample was limited; larger replications are needed, but the physiological rationale is anatomically sound.

Technique 2: ROCKING

Rocking replaces the in-out thrusting motion with a rolling, grinding, circular motion that maintains internal contact while creating sustained stimulation. Thrusting is optimised for the anatomy of the person doing the thrusting. Rocking is considerably more useful for the anatomy of the person receiving it.

The research is consistent: many women require 10 or more minutes of sustained clitoral stimulation to reach orgasm. Thrusting interrupts stimulation repeatedly. Rocking maintains that pressure continuously. Think of orgasm as a pressure system that needs to be built: thrusting is a pump with a slow leak; rocking seals the system.

Technique 3: SHALLOWING

Shallowing is the deliberate focus on the first 1 to 2 inches of the vaginal opening rather than pursuing depth. The vestibule and vaginal introitus contain the highest concentration of nerve endings in the entire vaginal region. The vestibular bulbs, part of the clitoral complex, flank this exact zone and are engorged during arousal. Shallowing is not withholding depth. It is targeting the most sensitive real estate in the area.

Technique 4: PAIRING

Pairing is simultaneous clitoral stimulation during penetrative sex. It is the simplest available adjustment and, statistically, the most universally effective. Given that 75 to 80% of women require direct clitoral stimulation to orgasm, pairing is not optional enhancement. For most women, it is a prerequisite.

Nobody put clitoral stimulation in the screenplay. This chapter is correcting the oversight.

Methods of Pairing

- Manual pairing by the receiving partner: the most common and most effective method.
- Manual pairing by the penetrating partner: requires communication about preferred pressure, location, and rhythm. Worth the conversation.

- Vibrator pairing: a small vibrator or clitoral suction device held against the clitoris during penetration.
- Position-based pairing: positions that naturally create clitoral contact through pubic-bone friction, such as woman-on-top with a forward lean or the CAT position.

Many women feel self-conscious about stimulating themselves during partnered sex, as though it communicates that their partner is insufficient. This discomfort is understandable, deeply culturally conditioned, and entirely worth overcoming. Direct clitoral stimulation during penetration is not a commentary on a partner's performance.

📊 THE CORE FOUR IN COMBINATION

- Angling + Pairing: anterior-wall stimulation combined with clitoral stimulation. Highest-success combination for penetrative orgasm.
- Rocking + Pairing: sustained pressure plus continuous clitoral stimulation. Replicates the pattern most associated with reliable female orgasm.
- Shallowing + Oral: external oral stimulation while shallow stimulation at the entrance creates dual-zone sensation.
- Full technique combination table: see Appendix B.

✦ CHAPTER SUMMARY — THE CORE FOUR

- Angling redirects penetration to the anterior wall where internal clitoral structures are located
- Rocking maintains sustained stimulation; thrusting interrupts it
- Shallowing targets the most richly innervated zone at the vaginal entrance
- Pairing adds direct clitoral stimulation — for most women, this is not optional
- The Core Four work best in combination; Angling + Pairing is the highest-success starting point

CHAPTER 5

Expanded Technique Toolkit

The Core Four in Chapter 4 are your foundation. This chapter is everything else: advanced positioning, edging, pelvic floor practice, breathwork, and the subtler adjustments that compound the basics into something genuinely extraordinary.

5.1 Advanced Positioning for Female Orgasm

Cowgirl (Woman on Top)

The receiving partner has full control of angle, depth, speed, and pressure. When the receiver leans forward 15 to 20 degrees, pubic bone contact with the clitoris becomes possible throughout. Add a vibrator and this position becomes one of the most reliable pathways to penetrative orgasm.

Modified Missionary with Pelvic Pillow

Elevating the receiver's hips by 10 to 15 cm with a pillow changes the angle of penetration toward the anterior wall. Combined with the CAT, this is one of the highest-success positions in the research literature.

Spooning

Deeply underrated. The receiver's hand has completely free access to the clitoris throughout, pairing is effortless, and the pace is naturally

more rocking than thrusting. Not glamorous. Very effective.

Flat Doggy (Prone Bone)

The receiver lies flat on their stomach. The mattress creates continuous pressure on the mons pubis and clitoris throughout. The angle naturally targets the anterior wall. A pillow under the hips increases the effect.

Scissors / Tribadism

Works for all anatomies. Full clitoral contact throughout. Natural rocking motion. Generously lubricated, this is one of the most direct routes to orgasm available.

5.2 The Edging Technique

Edging involves deliberately approaching but not reaching orgasm, reducing stimulation, and then rebuilding arousal. The cycle is repeated several times before allowing release. The eventual orgasm is significantly more intense when the nervous system has been repeatedly brought to the threshold and held there. Regular edging practice also builds body awareness and orgasmic reliability over time.

Practical instruction: when arousal reaches approximately 7 out of 10, reduce stimulation to around 3 out of 10 for 60 to 90 seconds. Rebuild. Repeat three to five times. Allow release on the final cycle.

5.3 Pelvic Floor Engagement

The pelvic floor muscles contract rhythmically during orgasm. This is the literal mechanism, not a metaphor. Practising voluntary pelvic floor contractions strengthens the muscles involved in orgasm and

can directly increase both intensity and frequency of orgasmic response.

The basic Kegel: contract the pelvic floor for 5 seconds, release for 5 seconds. Ten repetitions, three times daily. This is pleasure enhancement for women at every life stage, not just postpartum recovery work.

5.4 Breathwork and Orgasmic Breath

Breath is the overlooked master variable. Shallow breathing keeps the nervous system in a mild state of activation that actively inhibits deep arousal. Deep, rhythmic breathing activates the parasympathetic nervous system, which is the physiological state required for full sexual response.

Practise belly breathing during sex: consciously expand the abdomen on the inhale, release fully on the exhale. As arousal builds, breathe more audibly and slightly faster. This feeds oxygen to the building physiological response and signals to the body that it is safe to escalate.

★ EVIDENCE NOTE — BREATHWORK AND AROUSAL: MODERATE EVIDENCE

The parasympathetic link to breathwork is well-established in nervous system research. Direct controlled studies on breathwork and orgasm specifically are limited; the application draws on solid physiological foundations but has not yet been the subject of large RCTs in a sexual context.

✦ CHAPTER SUMMARY — EXPANDED TECHNIQUE TOOLKIT

• Position selection matters: cowgirl with forward lean and modified missionary are highest-evidence for penetrative orgasm
• Spooning and prone bone are underrated for natural clitoral contact

- Edging builds orgasmic intensity and reliability through threshold practice
- Pelvic floor training directly strengthens the muscles involved in orgasm
- Breathwork activates the parasympathetic state required for full arousal

CHAPTER 6

Oral Mastery & Manual Expertise

❧

For most women, the most reliable route to orgasm is not penetration. It is oral sex or skilled manual stimulation. This is well-documented, widely known to researchers, and somehow still underemphasised in mainstream sexual culture. This chapter corrects that.

6.1 Oral Sex: The Complete Guide to Cunnilingus

The Approach

Begin outside the genitals entirely. The inner thighs, the mons pubis, the crease where leg meets pelvis are richly innervated regions that respond to warm breath, light pressure, and slow attention. Beginning here builds anticipation and allows time for vasocongestion to begin. Jumping straight to the clitoris is like walking into a party before anyone has arrived.

Working Toward the Clitoris

Approach the clitoris indirectly before directly. Broad tongue strokes across the labia minora, pressure against the clitoral hood rather than directly on the glans, attention to the vestibule. The clitoral glans, once aroused, is extremely sensitive and often overstimulating at initial direct contact. The hood exists for a reason.

Rhythm and Consistency

This cannot be overstated: when something is working, do not change it. The most common error in oral sex is constant technique variation driven by performance anxiety or misread signals. When a woman is approaching orgasm, the arousal pattern is building through consistency. Changing technique at this point interrupts the build and requires starting again. When something is working, maintain it with conviction.

Incorporating Fingers

Adding one or two fingers, pad facing upward, 2 to 3 inches in, with a gentle come-hither motion toward the anterior wall, while maintaining oral stimulation creates simultaneous clitoral and G-spot area stimulation. For many women, this combination is significantly more effective than either alone. Do not add fingers without checking first.

ORAL SEX: DO'S AND DON'TS

- DO start slowly and build. The warmup is not optional.
- DO maintain rhythm when something is working. Consistency is the mechanism.
- DO stay present when she is close. This is not the moment to switch things up.
- DON'T rush to the clitoris before the surrounding tissue is warmed up.
- DON'T add fingers without checking first.
- DON'T stop abruptly when orgasm is close.
- DON'T treat it as a warmup for something else. It can be the main event.

REAL WOMEN, REAL STORIES — DANIELLE, 29, CANADA

"My previous partners always stopped right before I got there. Every time. I used to think something was wrong with my timing. Then I was with someone who just stayed. Who didn't need to change or escalate or perform. He just stayed. I have never experienced anything like it."

On what presence does that performance never can. Names and identifying details have been changed to protect privacy.

6.2 Manual Stimulation: The Hands as Instruments

External Technique

Lubrication is essential for external manual stimulation. A dry touch on the clitoral glans is uncomfortable in a way that does not build arousal. Use good-quality lubricant generously.

The most effective external manual techniques from the OMGYES research: circular motion with rhythmic circles over the clitoral hood; up-down stroke along the clitoral hood; figure-8 across the entire vulva; and tapping, light rapid taps directly on the glans that can trigger rapid escalation in women who respond to this technique.

Internal Manual Technique

G-spot stimulation requires adequate arousal first. The erectile tissue of the anterior wall is only engorged and accessible when the woman is genuinely aroused. Insert one finger, pad facing upward, 2 to 3 inches in. The G-spot area has a slightly different texture, sometimes described as ridged or spongy. Apply a come-hither curving motion toward the anterior wall.

The most effective combination: one or two fingers on the anterior wall while the thumb stimulates the clitoris simultaneously. This is the manual equivalent of the blended orgasm. It takes coordination. It is worth it.

✦ CHAPTER SUMMARY — ORAL MASTERY & MANUAL EXPERTISE

• For most women, oral and manual stimulation are more reliable routes to orgasm than penetration
• Begin away from the genitals; vasocongestion takes time
• When something is working, maintain it — consistency builds orgasm, variation breaks it
• Lubrication is essential for manual stimulation
• The manual blended technique — anterior wall plus clitoral stimulation — is among the most effective combinations available

CHAPTER 7

Solo Practice as the Foundation of Partnered Pleasure

❧

Let us talk about masturbation. Not as something to confess or justify or apologise for. As something useful.

Solo sexual practice is the best available research method for learning about your own body. It is private, available, and requires no coordination with another person's schedule. It is the laboratory for self-knowledge, and self-knowledge is the single most powerful tool available for improving partnered sex.

★ EVIDENCE NOTE — SOLO PRACTICE AND PARTNERED SATISFACTION: STRONG EVIDENCE

Women who masturbate regularly report higher rates of orgasm during partnered sex, higher sexual self-esteem, and greater communication quality with partners (Herbenick et al., 2010; OMGYES 2016). Vibrator use is associated with higher overall sexual satisfaction, not lower. Self-knowledge of one's own pleasure responses is among the strongest predictors of partner communication quality.

7.1 The Guided Self-Exploration Framework

Stage 1: The Body Audit (Week 1)

Before introducing orgasm as a goal, spend time simply touching and mapping your body without expectation. Not just the genitalia. The neck, the inner arm, the collarbone, the lower back, the inner thigh.

Discover what kinds of touch feel good. This is information-gathering, not performance.

Stage 2: Genital Mapping (Week 2)

Using a mirror and good lighting, explore the vulva visually first. Identify the structures from Chapter 2. Then begin touching each area and noting sensation. There is no correct response. This is cartography, not evaluation.

Stage 3: Arousal Building (Weeks 3 to 4)

Introduce intentional arousal: fantasy, erotic reading, or audio. Begin clitoral stimulation using fingers or a vibrator. Focus on what builds sensation rather than on reaching orgasm. Curiosity is more useful than urgency here.

Stage 4: Partner Translation (Ongoing)

Once you have a reliable map of your own responses, practise translating that knowledge to a partner. This can be verbal, physical through guiding their hand, or demonstrative. The knowledge is not useful only to you. It is the most valuable thing you can bring to a partnership.

✦ CHAPTER SUMMARY — SOLO PRACTICE

- Solo practice is not supplemental — it is the primary laboratory for sexual self-knowledge
- Women who masturbate regularly are significantly more likely to orgasm with a partner
- The four-stage framework builds from body awareness to genital mapping to arousal building to partner translation
- Vibrator use correlates with better sexual function, not decreased sensitivity

PART THREE

THE MIND-BODY CONNECTION

Because the Brain Is the Largest Sex Organ

CHAPTER 8

The Psychology of Female Orgasm

❧

Everything we have covered so far is necessary but not sufficient. Because the female orgasm does not happen only in the body. It happens in the brain first.

★ **EVIDENCE NOTE — BRAIN IMAGING AND ORGASM: STRONG EVIDENCE**

Neuroscientists using fMRI have found that a full orgasm involves nearly every region of the brain (Komisaruk et al., 2011). This whole-brain activation is closer to a mild altered state than a simple physical reflex. The psychological conditions under which it becomes possible matter enormously.

8.1 The Dual Control Model

The most useful framework for understanding female sexual response comes from research by John Bancroft and Erick Janssen: the Dual Control Model. It proposes that sexual response is governed by two competing systems operating simultaneously.

The Sexual Excitation System (SES) — the accelerator — responds to sexually relevant stimuli and sends signals that say go. The Sexual Inhibition System (SIS) — the brakes — responds to potential threats, distractions, and inhibitory cues and sends signals that say stop. Sexual response is the net result of acceleration minus braking.

The critical finding: on average, women's brakes tend to be more sensitive and more numerous than men's. This does not mean women

are less sexual. It means that for many women, removing inhibitory signals is more important than adding excitatory ones. You can press the accelerator all you like, but if the handbrake is on, you are not going anywhere.

COMMON BRAKE FACTORS FOR WOMEN

- Body image concerns and self-consciousness during sex
- Worry about partner's pleasure, satisfaction, or judgment
- Unresolved relationship tension or emotional distance
- Anxiety about pregnancy, STIs, or being overheard
- Stress, fatigue, or mental preoccupation with unrelated concerns
- Past negative sexual experiences or trauma
- Performance anxiety, including anxiety specifically about not orgasming
- Cultural or religious shame around sexuality

8.2 Spectatoring

In the 1970s, Masters and Johnson described spectatoring: the habit of mentally stepping outside one's own body during sex to observe and evaluate oneself from an imagined external perspective. It sounds like this: How do I look from this angle? Am I taking too long? Is my partner bored?

This internal monologue is among the most effective orgasm inhibitors available without prescription. When attention is directed outward toward evaluation, it is not available for the physical experience unfolding in the body.

The solution is not willpower. It is practice. Mindfulness-based interventions, sensate focus exercises, and breath anchoring are all evidence-based tools for reducing spectatoring.

8.3 Mindfulness and Embodiment Practices

• Body scan before sex: spend 5 minutes in a simple body scan beginning at the feet and moving slowly upward, noticing sensation without judgment. This is a nervous system protocol for arriving in your body rather than in your head.
• Sensate focus exercises: developed by Masters and Johnson and refined by sex therapists since, these are structured touch sessions with the explicit goal of noticing sensation without performance pressure. Research shows significant improvements in orgasm frequency and sexual satisfaction.
• The permission practice: many women have an unconscious conditional structure around orgasm. The permission practice is noticing this condition and choosing, deliberately, to remove it. Orgasm is not something to be earned.

REAL WOMEN, REAL STORIES — YUNA, 41, SOUTH KOREA

"I had spent years in my own head during sex. Monitoring. Managing. Making sure it was good for him. My therapist called it spectatoring and I remember thinking, that's exactly it. The shift came slowly. I started using breath to bring myself back into my body. It took about six months of practice before something actually changed. But it did change."

On returning to yourself, one breath at a time. Names and identifying details have been changed to protect privacy.

✦ CHAPTER SUMMARY — THE PSYCHOLOGY OF FEMALE ORGASM

• Orgasm is a whole-brain event — psychology is not separate from physiology, it is physiology
• The Dual Control Model: for many women, removing brakes matters more than pressing the accelerator
• Spectatoring is among the most effective orgasm inhibitors — and it responds to mindfulness practice
• Sensate focus and body scan practices have evidence bases for improving orgasm frequency
• Orgasm does not need to be earned. It is a physiological response that deserves space.

CHAPTER 9

Emotional Safety, Trust & Vulnerability

There is a physiological reason why emotionally safe sex is better sex. Not just more enjoyable, not just more comfortable, but physiologically different in its capacity for orgasmic response.

9.1 The Nervous System Case for Emotional Safety

The human nervous system cannot simultaneously be in a state of vigilance and a state of deep physical surrender. These are mutually exclusive modes. Orgasm is a parasympathetic event. It requires, at its neurological core, a felt sense of safety.

When the nervous system perceives threat — not just physical threat but social threat, the subtle sense that one is being judged or not fully seen — it activates a cascade of physiological responses. Heart rate increases slightly. Muscle tension rises. The genitals receive less of the vasocongestion that arousal requires. This is not psychological weakness. It is physiology.

9.2 The Vulnerability Paradox

Orgasm requires a degree of physical and emotional surrender that most women have been culturally taught to resist. Orgasm involves

losing a degree of control over the body: involuntary sounds, involuntary muscle movements, a brief altered state. It is quite literally a moment of being out of control in front of another person.

The paradox: the same culture that produced the orgasm gap has also taught women to maintain composure, monitor their appearance, manage others' comfort, and perform rather than experience. These are precisely the habits that prevent orgasm. We are asking women to be fully vulnerable in their most intimate moments while simultaneously equipping them with every tool for staying defended.

The pathway is not to force vulnerability. It is to build it gradually, through communication, emotional intimacy, and incremental trust. Real safety is earned, not assumed. Build the architecture first.

9.3 Practical Pathways to Emotional Safety

Repair Before Intimacy

Attempting sex when significant emotional tension is present rarely ends well. A repair conversation, even an incomplete one, matters.

The Decompression Transition

Create a ritual transition between the rest of the day and the intimate space: a shower, a conversation, a shared meal, or 20 minutes of deliberate non-sexual physical affection. The nervous system needs to downregulate from daily stress before it is available for full arousal. This is not foreplay. It is pre-foreplay.

Verbal Reassurance During Sex

Simple, genuine verbal presence from a partner — I'm here, take your time — can significantly reduce spectatoring and increase the felt sense of safety. Many women have never experienced this. Many

partners have never offered it. It is not complicated. It is extraordinarily effective.

✦ CHAPTER SUMMARY — EMOTIONAL SAFETY, TRUST & VULNERABILITY

• Orgasm is a parasympathetic event — it requires felt safety at the neurological level
• Social threat activates the same protective response as physical threat
• Vulnerability cannot be forced; it can only be built incrementally through trust
• Repair before intimacy, decompression rituals, and verbal reassurance are practical tools
• Emotional presence from a partner is more predictive of orgasm than technical skill

CHAPTER 10

Desire, Libido & the Responsive Model

Here is a question that has led many women into unnecessary misery: what is wrong with my libido? The question assumes a model of desire that works like hunger: it arises spontaneously, independent of context, before stimulation rather than in response to it. This model is wrong for a large proportion of women.

10.1 Spontaneous vs. Responsive Desire

Spontaneous desire arises without obvious context or stimulation. Responsive desire emerges in response to stimulation or erotic context. It does not show up before the encounter. It shows up during it, once conditions have been created that allow it to develop. This is the normal baseline desire pattern for many women, particularly in longer-term relationships.

The critical practical implication: if your desire is primarily responsive, waiting until you feel like it before initiating sex is a category error. You will not feel like it until the conditions exist to create the feeling.

★ **EVIDENCE NOTE — RESPONSIVE DESIRE: STRONG EVIDENCE**

Basson's circular model of female sexual response (2002, Journal of Sex & Marital Therapy) established responsive desire as a normal and common pattern,

particularly in long-term relationships. Emily Nagoski's synthesis in Come As You Are (2015) popularised this finding for general audiences. It is now considered a cornerstone of sex-positive clinical practice.

SPONTANEOUS VS. RESPONSIVE DESIRE

• Spontaneous desire: arrives before any sexual stimulation, apparently from nowhere
• Responsive desire: emerges during stimulation, not before it
• Neither is more real or more healthy than the other
• Responsive desire is frequently misread as low libido — it is not
• Many women discover responsive desire is their baseline only after reading about it for the first time

10.2 When Libido Genuinely Decreases

Responsive desire is not the same as low libido. Genuine reductions in libido have identifiable causes: hormonal shifts around perimenopause and menopause, certain medications especially SSRIs and combined oral contraceptives, thyroid dysfunction, chronic fatigue, depression, significant relationship distress, or unprocessed trauma. These are real and addressable. The appropriate response is investigation, not self-blame.

★ EVIDENCE NOTE — MEDICATIONS AND LIBIDO: STRONG EVIDENCE

SSRIs are among the most commonly prescribed medications worldwide and have a well-documented effect on sexual desire and orgasm capacity. An estimated 30 to 70% of users experience some sexual side effect (Clayton et al., 2014). If you are taking an SSRI and have noticed changes in desire or orgasm, discuss it with your prescriber. Alternatives exist. You should not have to choose between mental health and sexual health.

✦ CHAPTER SUMMARY — DESIRE, LIBIDO & THE RESPONSIVE MODEL

• Responsive desire is normal, common, and not a sign of low libido

• For responsive desire, creating conditions matters more than waiting to feel ready
• Genuine libido decreases have identifiable causes: hormones, medications, stress, trauma
• SSRIs affect libido in 30–70% of users — this is medical, not imagined
• Investigation, not self-blame, is the correct response to genuine desire changes

CHAPTER 11

Communication as Foreplay

❧

The research is unambiguous: communication is the single highest-leverage intervention available for improving sexual satisfaction. More effective than any position, more impactful than any technique. And it is also the most consistently avoided.

★ EVIDENCE NOTE — COMMUNICATION AND SEXUAL SATISFACTION: STRONG EVIDENCE

Mark et al. (2014, Archives of Sexual Behavior) found that sexual communication was among the strongest predictors of sexual satisfaction and orgasm frequency in both short- and long-term relationships. Vannier & O'Sullivan (2012) found communication quality outperformed frequency, novelty, and technique in predicting satisfaction. This finding is among the most replicated in sexual science.

11.1 Why We Don't Talk About It

Sexual communication is awkward for reasons that are entirely understandable and not at all inevitable. We were not taught how. The script says desire should be wordless and intuitive. Many women were socialised to prioritise a partner's comfort over their own needs. And there is the very reasonable fear that saying that's not quite right will be heard as you're doing it wrong.

All of these are real. None of them are insurmountable. And the cost of not communicating, measured in years of unsatisfying sex and

widening emotional distance, is far higher than the awkwardness of the conversation.

11.2 The Three Windows

Window 1: Before

The desire conversation. What do you want? What are you curious about? What felt wonderful last time? This conversation happens outside the sexual context entirely — over dinner, on a walk, in a text message. Low-stakes and exploratory. No performance pressure because nothing is happening right now.

Window 2: During

Real-time guidance. Right there. A little softer. Don't stop. This should feel like giving your partner a gift rather than evaluating their performance.

Window 3: After

The debrief. What I loved most about that was... Next time, I'd love to try... This window is the most underused of the three and one of the most powerful. A post-encounter conversation reinforces what worked, builds knowledge for the future, and creates a shared language around pleasure that compounds over time.

SAMPLE COMMUNICATION SCRIPTS

Opening Before: 'I've been thinking about what really feels good for me lately, and I'd love to share some of that with you. Can we find a time to talk about it?'

In the Moment (positive): 'Yes, exactly like that. Please don't change anything.'

In the Moment (redirecting): 'Can you try it a little higher?' Frame guidance as information, not criticism.

After Sex: 'What I loved most was when you... I'd love more of that.'

The Big Conversation: 'I want to be honest with you about something that matters to me. I care about our sex life and I'd love for us to be more open about what we both enjoy.'

✦ CHAPTER SUMMARY — COMMUNICATION AS FOREPLAY

• Communication is the single highest-leverage intervention for sexual satisfaction — above technique, position, or duration
• The three windows: Before (desire conversation), During (real-time guidance), After (debrief)
• Scripts reduce the awkwardness of first conversations
• Frame guidance as a gift, not criticism
• The debrief window is the most underused and one of the most powerful

PART FOUR

THE EROGENOUS MAP

Spots, Zones & the Full Geography of Pleasure

CHAPTER 12

G-Spot, A-Spot, C-Complex & Beyond

The internal female anatomy contains multiple distinct regions capable of producing profoundly different orgasmic experiences. A consistent theme across all internal exploration: arousal first, always. The structures described in this chapter are all either inaccessible or actively uncomfortable when a woman is not fully aroused.

12.1 The G-Spot: What It Actually Is

The term G-spot was popularised by Beverly Whipple in the 1980s, named after German gynaecologist Ernst Grafenberg who described the area in 1950. The debate about whether it exists as a distinct structure has been, from an anatomical perspective, somewhat beside the point.

The anterior vaginal wall at approximately 2 to 3 inches depth overlies the urethral sponge and the internal clitoral structures. When a woman is fully aroused, this tissue becomes engorged and takes on a slightly different texture, often described as ridged or spongy. This is the CUV (clitorourethrovaginal) complex, and stimulation of this area produces a distinct and often intense orgasmic response in a significant proportion of women.

★ **EVIDENCE NOTE — WHERE THE SCIENCE STANDS ON THE G-SPOT: CONTESTED**

Whether the G-spot is a distinct anatomical structure or simply the internal clitoris accessed through the anterior vaginal wall is genuinely contested among researchers. Helen O'Connell's 1998 anatomy paper and subsequent ultrasound studies (Buisson et al., 2021, Clinical Anatomy) strongly support the latter — that what we call the G-spot is the internal clitoris. However, other researchers maintain that the CUV complex constitutes its own functionally distinct zone. Both sides agree that anterior-wall stimulation produces significant responses in many women. The naming debate should not distract from the technique.

How to Find It

Insert one or two fingers, pad facing upward toward the front of the body, 2 to 3 inches in. Make a come-hither curving motion against the anterior wall. When she is sufficiently aroused, you will feel a change in texture. Apply sustained, rhythmic pressure there.

12.2 The A-Spot: The Underknown Zone

The A-spot, or Anterior Fornix Erogenous zone, is located at 4 to 6 inches depth on the anterior wall, in the fold between the cervix and the anterior vaginal wall. First described by Malaysian sexologist Chua Chee Ann in the 1990s.

This zone requires both depth and full arousal to be accessible. With sufficient arousal, stimulation of the A-spot frequently produces an immediate increase in vaginal lubrication and a profound, deep arousal that many women describe as qualitatively different from clitoral stimulation.

★ **EVIDENCE NOTE — A-SPOT EVIDENCE: MODERATE**

Chua Chee Ann's 1997 clinical study found compelling results for A-spot stimulation, including rapid lubrication increase in women who reported previous dryness. The

study was methodologically limited in sample size. Independent large-scale replications have not yet been published. The anatomical rationale is sound but treat outcomes as individual rather than universal.

12.3 The Cervix and Deep Stimulation

The cervix sits at the apex of the vaginal canal. Its sensitivity varies enormously: some women find cervical contact exquisitely pleasurable; others find it uncomfortable or painful. Cervical orgasm is possible for some women and is described as a transcendent, full-body, sometimes emotionally charged experience. It is only accessible at peak arousal and is not available to all women regardless of how much they want it to be. If deep stimulation is painful rather than pleasurable, this is physiological information, not failure.

📊 THE GOLDEN RULE OF INTERNAL EXPLORATION

• Arousal first. Always. This is non-negotiable.
• The internal erectile tissue becomes engorged and accessible only with full arousal.
• Recommended minimum: 15 minutes of external stimulation before any internal exploration.
• If it hurts, stop and recalibrate. Pain is physiological information, not a challenge to overcome.
• This is information-seeking, not goal-achievement. Treat it that way.

✦ CHAPTER SUMMARY — G-SPOT, A-SPOT, C-COMPLEX & BEYOND

• The G-spot is most accurately understood as the internal clitoris accessed through the anterior wall — though the anatomical debate continues
• Both major research positions agree: anterior-wall stimulation works for many women
• The A-spot is located deeper (4–6 inches) and requires full arousal to access
• Cervical sensitivity varies enormously — for some women it is transcendent, for others uncomfortable
• Arousal first, always: without it, internal zones are inaccessible or painful

CHAPTER 13

The Clitoris: The Complete Architecture

❧

We introduced the clitoris in Chapter 2. But the clitoris deserves its own chapter — not because the information is entirely new at this stage, but because bringing it to the centre of the conversation is itself the point. It is, for the majority of women, the primary pleasure organ, and it has been, for most of recorded history, the most misunderstood structure in human anatomy.

13.1 Complete Clitoral Anatomy

The Glans Clitoridis

The visible external tip, approximately 1 cm when unaroused and up to 3 cm when fully engorged. Contains approximately 8,000 nerve endings in an area smaller than a fingertip. The nerve density is double that of the entire penis.

The Clitoral Hood (Prepuce)

A fold of skin that protects the glans. Whether stimulation through the hood or direct contact is preferred is highly individual and worth exploring early.

The Clitoral Body

Extends approximately 2 to 3 cm internally from the glans toward the pubic bone, beneath the skin of the mons pubis.

The Clitoral Crura

Two arms that extend from the body along the pubic arch, each 5 to 9 cm in length. Entirely internal. They surround the vaginal canal and respond to penetrative stimulation.

The Vestibular Bulbs

Two teardrop-shaped erectile bodies, one on each side of the vaginal opening. They become engorged with blood during arousal and are directly stimulated by pressure at the vaginal entrance.

13.2 The Clitoral-Vaginal Connection Explained

The persistent question — is vaginal orgasm different from clitoral orgasm? — is largely answered by anatomy. The clitoral crura and vestibular bulbs surround and respond to the vaginal canal. What is experienced as a vaginal orgasm is, in most cases, clitoral stimulation mediated through the internal structures.

There is no established anatomical mechanism by which the vaginal walls alone could produce orgasm in the majority of women. The vaginal canal is not particularly richly innervated past its first two inches. This does not mean that all orgasms feel the same. They do not. The difference is in how the clitoral structure is being stimulated, not in whether it is involved.

★ EVIDENCE NOTE — THE VAGINAL-CLITORAL QUESTION: STRONG EVIDENCE FOR CLITORAL INVOLVEMENT

O'Connell et al. (1998, Journal of Urology) established the full internal architecture of the clitoris. Buisson et al. (2021) used ultrasound imaging to show that penetration stimulates the internal clitoral structures. While 'all orgasms are clitoral' is an oversimplification that some researchers contest at the neuroscience level, clitoral involvement in all orgasms is well-supported anatomically. The debate sits primarily in neurological framing, not anatomy.

13.3 How the Techniques Connect

Having now established the complete architecture, every technique from earlier chapters makes fuller sense. Angling directs penetration toward the internal crura and vestibular bulbs. Shallowing targets the vestibular bulbs directly at the entrance. Rocking maintains sustained pressure on the internal structure. Pairing adds direct external stimulation to the already-stimulated internal structure.

Every technique in this book is ultimately in service of stimulating the clitoris. The pathway to female orgasm has always been there. We have simply been calling it by the wrong names and approaching it from the wrong directions.

✦ CHAPTER SUMMARY — THE CLITORIS: THE COMPLETE ARCHITECTURE

- The clitoris is a multi-part internal organ; the visible glans is only 10% of the structure
- The crura and vestibular bulbs surround the vaginal canal and respond to penetration
- All orgasms involve clitoral activation, direct or indirect
- Every technique in this book targets the clitoris through different pathways
- Understanding the full architecture explains every technique and why each works

CHAPTER 14

Cervical, Blended & Full-Body Orgasms

❧

Orgasm is not a single thing. That may be the most useful sentence in this chapter.

14.1 The Orgasm Spectrum

Clitoral Orgasm

The most common, most reliably triggered, and most rapidly achieved orgasm available. It originates from direct stimulation of the clitoral glans and is characterised by a sharp, intense, localised sensation. This is the baseline.

G-Spot Orgasm

Triggered by sustained anterior-wall stimulation. Builds more slowly, often with a sensation of pressure or fullness. When it arrives, it tends to be deeper, more spread through the pelvis, and often accompanied by a sense of internal release.

Blended Orgasm

Occurs with simultaneous clitoral and G-spot stimulation. Combines the intensity of clitoral orgasm with the depth of G-spot orgasm and tends to be longer-lasting than either alone.

A-Spot Orgasm

Triggered by deep anterior wall stimulation near the cervix. Less commonly discussed but described by women who have experienced it as profound and spreading.

Cervical Orgasm

For women who can access it, described as transcendent, full-body, sometimes emotionally cathartic, and qualitatively different from any other orgasm type. Only possible at very high arousal levels. Not all women experience this and that is not a shortcoming.

Full-Body Orgasm

A wave-like orgasmic sensation that moves through the entire body rather than being localised in the genitals. Associated with breathwork practices and extremely high arousal states.

Multiple Orgasms

Sequential orgasms occurring without a full refractory period. The capacity can be developed through practice, edging, and pelvic floor work.

★ EVIDENCE NOTE — ORGASM TYPES: VARIED EVIDENCE LEVELS

Clitoral orgasm is strongly evidenced and well-characterised. Blended and G-spot orgasms are moderately evidenced with consistent clinical reporting but limited large controlled studies. Cervical orgasm is documented primarily through case studies and qualitative research (Komisaruk & Whipple, 2011); no large-scale RCTs exist. Full-body orgasm associated with breathwork is documented clinically but lacks neuroimaging studies. Consistency of experience reports does not equal the same evidentiary standard as replicated controlled studies — these distinctions matter for calibrated expectations.

14.2 Sleep and Exercise-Induced Orgasms

Sleep orgasms occur in approximately 37% of women and are triggered by erotic dreams and nocturnal genital engorgement. They are entirely normal. Exercise-induced orgasms, sometimes called coregasms, occur during physical exercise involving abdominal and pelvic floor engagement. Approximately 10 to 15% of women report having experienced this.

✦ CHAPTER SUMMARY — CERVICAL, BLENDED & FULL-BODY ORGASMS

- Orgasm is a spectrum, not a single experience
- Clitoral orgasm: the most common and reliably achieved; the baseline
- Blended orgasm combines clitoral and G-spot stimulation for longer, deeper experience
- Cervical and full-body orgasms are real but require high arousal and are not universal
- Evidence levels vary by type: calibrate expectations accordingly

PART FIVE

BODIES IN FULL

Trauma, Disability, Neurodivergence & What Partners Carry Too

CHAPTER 15

When the Body Has a History: Trauma, Healing & Sexual Recovery

❧

For many women, the body has a history. Sometimes that history is quiet — a vague sense of not feeling fully present during sex, a reflex to go somewhere else in the mind, a tension that never quite releases. Sometimes that history is louder: the aftermath of assault, abuse, neglect, or a medical experience that did not ask for consent.

This chapter is not a clinical trauma treatment guide. It is written for women who already know that their sexual experience is shaped by things that happened before this relationship, before this encounter, before this moment. It is written in the belief that healing is possible, that sexual pleasure is available even in the context of trauma history, and that the path there requires knowledge, patience, and self-compassion rather than willpower.

15.1 How Trauma Affects Sexual Response

Trauma — understood broadly as any experience that overwhelmed the nervous system's capacity to cope — leaves residue. In the context of sexual experience, this residue most commonly appears as: hyperarousal, a state of heightened vigilance that keeps the nervous system from fully relaxing into arousal; dissociation, the sense of floating above or outside the body during sex; arousal-inhibition

patterns; and trigger responses, specific sensory experiences that activate trauma memories even in otherwise safe contexts.

These are not character flaws. They are adaptive responses that worked at one time, in conditions of genuine danger, and are now activating in the wrong context. The nervous system cannot always tell the difference between then and now.

> ★ **EVIDENCE NOTE — TRAUMA AND THE BODY: STRONG EVIDENCE**
>
> Van der Kolk's foundational research (The Body Keeps the Score, 2014) and Levine's somatic experiencing work have established that trauma is stored somatically, in the body's tissues and nervous system patterns. Brain imaging studies show traumatic memories activate survival systems in ways that override the rational brain's knowledge that danger has passed. This is not metaphorical. It is neurological.

15.2 Trauma-Informed Approaches to Sexual Healing

The Safety Ladder

Sexual healing for trauma survivors is not about pushing through discomfort to reclaim sexuality. It is about building a ladder of felt safety, one rung at a time, until the nervous system genuinely recognises the current context as different from the past.

- Non-sexual touch first: the back, the hands, the feet.
- Gradual expansion of touch as each new boundary is met with safety and consent.
- Explicit permission to stop at any point without explanation.
- Explicit communication with a partner about what is helpful and what activates a threat response.

Grounding Techniques During Sex

When dissociation or a trigger response begins during sex: name five things you can see in the room. Feel the weight of your body against the bed. Take a deliberate, audible breath. Speak a word or signal agreed with your partner that means pause. These are not interruptions of intimacy. They are the practice of staying present.

Working with a Therapist

Trauma-informed sex therapy, somatic experiencing, EMDR, and Sensorimotor Psychotherapy all have evidence bases for helping trauma survivors reclaim sexual wellbeing. This is not weakness. It is precision care for a specific kind of wound.

REAL WOMEN, REAL STORIES — AMARA, 36, NIGERIA

"I was assaulted at 22. For years afterward, I went through the motions. I got good at performing presence while being completely absent. When I finally told my partner what was happening inside me, he cried. Not for himself. For me. That was the beginning. He started asking me, what do you need right now, before anything else. Sometimes the answer was just hold still. Don't move. Let me feel that this is safe. We went slowly. We are still going slowly. But I am actually here now."

On what healing looks like when someone chooses to stay. Names and identifying details have been changed to protect privacy.

15.3 For Partners of Trauma Survivors

Your partner's responses — the flinching, the going quiet, the sudden shift in energy, the need to stop — are not rejections of you. They are the nervous system doing the only thing it knows how to do, which is protect. Your role is not to fix this or accelerate it. Your role is to be reliably, consistently, boringly safe.

Ask, don't assume. Check in genuinely and without agenda. Accept a no without making it about you. And do your own work. A partner's trauma history sometimes activates unresolved material in the other person too. There is no shame in that. There is just the work.

📊 ORGANISATIONS THAT CAN HELP

• RAINN (Rape, Abuse & Incest National Network): rainn.org — 24/7 hotline: 1-800-656-HOPE (US)
• AASECT (American Association of Sexuality Educators, Counselors and Therapists): aasect.org
• International Society for Traumatic Stress Studies (ISTSS): istss.org
• One Billion Rising: onebillionrising.org — global anti-violence network with country-specific resources
• If you are outside the US, search for a certified trauma-informed therapist in your country using the ISTSS directory

✦ CHAPTER SUMMARY — TRAUMA, HEALING & SEXUAL RECOVERY

• Trauma is stored somatically — in the nervous system, not just in memory
• Hyperarousal, dissociation, and trigger responses are adaptations, not character flaws
• The safety ladder builds felt safety incrementally; it is not a fast process and is not supposed to be
• Grounding techniques and agreed pause signals keep healing practical during intimacy
• Partners: your role is reliable safety, not fixing or accelerating

CHAPTER 16

Every Body: Disability, Chronic Illness & Neurodivergence

The majority of books on female sexuality are written for a body that does not have chronic pain, does not take eight medications, has not had a mastectomy, does not live with sensory processing differences, and does not use a wheelchair. This chapter acknowledges explicitly that the principles of this book apply in modified and creative forms to a much wider range of bodies than the mainstream sexual script acknowledges.

16.1 Chronic Pain and Pelvic Pain Disorders

Conditions including endometriosis, vaginismus, vulvodynia, interstitial cystitis, and pelvic floor dysfunction affect a significant proportion of women and directly impact sexual experience. The orgasm gap for women with these conditions is not simply an education problem. It is a medical problem, a communication problem, and an emotional problem simultaneously.

Vaginismus is often misdiagnosed or dismissed for years. It is treatable, typically through a combination of pelvic floor physiotherapy, graduated dilator use, and psychological support. Vulvodynia responds to a range of medical and physiotherapy interventions. Neither of these conditions means sexual pleasure is

unavailable. It means the path there requires more knowledge and more support.

★ **EVIDENCE NOTE — PELVIC PAIN: STRONG EVIDENCE FOR TREATMENT**

Pelvic floor physiotherapy has strong evidence for improving pelvic pain conditions including vaginismus and vulvodynia (Rosenbaum & Owens, 2008; Bergeron et al., 2015). It is significantly underutilised globally. If penetration is consistently painful, a pelvic floor physiotherapist should be part of your healthcare team. Do not accept 'it's just how you are' as a medical opinion.

💬 **REAL WOMEN, REAL STORIES — SOFIA, 31, SPAIN**

"I was 27 before anyone told me what it was. Before that, I had been told by three different doctors that I was tense, that I needed to relax, that maybe I wasn't ready. I was ready. My body had a medical condition and nobody had bothered to name it. When I finally saw a pelvic floor physio, she looked at me on the first appointment and said: This is treatable. Those two words changed everything."

On what it means to finally be believed. Names and identifying details have been changed to protect privacy.

16.2 Disability and Sexuality

Disability does not end sexual desire. It changes the logistics. For women with physical disabilities, this may mean modifications to positioning, the use of adaptive equipment, or redefining what sex means and what counts as satisfying. The most consistent finding in research on disability and sexuality is this: women who receive information tailored to their condition report significantly higher sexual satisfaction than those who do not. Information is not a luxury. It is the intervention.

16.3 Neurodivergence and Sexual Experience

Autistic women, and women with ADHD, sensory processing differences, or anxiety disorders, often have sexual experiences that differ significantly from neurotypical norms in ways that are rarely discussed. Sensory hypersensitivity can make certain types of touch intensely uncomfortable rather than pleasurable. The most useful shift for neurodivergent women: explicit over implicit. Ask directly and answer directly. Negotiate sensory preferences openly and without embarrassment. Different is not broken.

16.4 Cancer, Surgery & Altered Bodies

Mastectomy, hysterectomy, oophorectomy, and cancer treatments alter the body in ways that can profoundly affect sexual experience and identity. Nerve changes, hormonal shifts, changes in erogenous zone sensitivity, and body image adjustments are real and frequently left unaddressed by medical teams who are understandably focused on survival outcomes. Sexual health after cancer treatment deserves dedicated attention. An increasing number of oncology programmes now include sexual health specialists. Ask for a referral if one is not offered.

✦ CHAPTER SUMMARY — EVERY BODY: DISABILITY, CHRONIC ILLNESS & NEURODIVERGENCE

- Disability and chronic illness change the logistics of pleasure, not the legitimacy of the desire
- Pelvic pain conditions are medical, not psychological — and they are treatable
- Pelvic floor physiotherapy is underutilised and has strong evidence
- For neurodivergent women: explicit communication is not a workaround, it is the tool
- Cancer and surgical changes deserve dedicated sexual health support — ask for it

CHAPTER 17

The Partner's Chapter: What He Carries Too

This book is primarily written for women. But the orgasm gap is not a problem that women create alone or solve alone. Male partners are inside it too, shaped by it, often burdened by it in ways they rarely have a language for. If you are a male partner reading this book, or reading this chapter first, welcome. You are exactly where you should be.

17.1 What Men Inherit

Men do not arrive in the bedroom from nowhere. They arrive carrying everything they were taught, or not taught, about sex. And what most men were taught is this: your job is to perform, your performance is measured by her orgasm, and if she doesn't orgasm, you failed. This is an extraordinarily high-pressure script that produces exactly the kind of performance anxiety, rushing, and technique-switching that we have spent this entire book explaining is counterproductive.

17.2 The Performance Anxiety Loop

Here is how the loop typically works. He tries a technique. It does not immediately produce the response he expects. He interprets the absence of visible response as failure. Anxiety increases. He either

switches techniques rapidly — introducing the inconsistency that Chapter 6 explains is one of the primary barriers to female orgasm — or he escalates stimulation past the point of her comfort.

She, sensing his anxiety, begins managing his feelings rather than attending to her own experience. She performs. He is temporarily reassured. Both of them are now further from genuine connection and further from her orgasm than they were five minutes ago.

The solution is not trying harder. It is trying differently. Specifically: staying present, maintaining consistency when something is working, and shifting the definition of success from her orgasm to her experience.

★ EVIDENCE NOTE — WHAT RESEARCH SAYS ABOUT MALE PARTNER BEHAVIOUR: STRONG EVIDENCE

Multiple studies have found that women with partners who are curious, attentive, and communicative report dramatically higher orgasm rates than women with technically skilled but emotionally inattentive partners (Frederick et al., 2018; Kleinplatz et al., 2009). Presence outperforms expertise. This is among the most consistent and practically useful findings in sexual satisfaction research.

17.3 Erectile Concerns and the Cycle of Avoidance

Erectile dysfunction affects a significant proportion of men, particularly over 40, and its psychological impact on a partnership's sexual life is profound. When a man experiences erectile difficulty, shame frequently follows. Shame produces avoidance. Avoidance reduces intimacy, which increases anxiety, which makes erectile function worse.

Erectile dysfunction has well-documented medical causes including cardiovascular disease, diabetes, medication side effects, and low testosterone. It is treatable. It is not a verdict on the relationship. And it is not something a partner caused.

17.4 Sexual Conditioning in Men

Male sexuality is shaped by pornography consumption at a scale that has no historical equivalent. Studies consistently find that heavy consumption is associated with higher rates of sexual dissatisfaction, unrealistic expectations about female sexual response, and lower ability to read real partner cues. This is not a moral judgment. It is a functional observation about what extended exposure to a highly distorted sexual script does to learned expectations.

This book is, among other things, a different manual. Share it.

REAL WOMEN, REAL STORIES — LENA, 43, SOUTH AFRICA

"He said it once, very quietly: I always felt like I was supposed to know. Like asking would mean I didn't know. And not knowing meant I wasn't good enough. I had never thought about it from that side before. We had been together for nine years and I had no idea he was carrying that. I had been faking. He had been pretending he knew. We had both been protecting each other from a conversation that would have changed everything."

On the silence that costs both people. Names and identifying details have been changed to protect privacy.

CHAPTER SUMMARY — THE PARTNER'S CHAPTER

- Men arrive carrying a performance script that actively works against female orgasm
- The performance anxiety loop: his anxiety creates her performance, which removes both from genuine connection

• Presence outperforms expertise in every study that has asked this question
• Erectile concerns are medical, treatable, and not a verdict on the relationship
• The most useful thing a partner can bring is curiosity, consistency, and the willingness to ask

PART SIX

CONTEXT & CULTURE

Pleasure Across the Lifespan & Closing the Gap Together

CHAPTER 18

Pleasure Across the Lifespan

Female sexuality is not a fixed point. It is a landscape that shifts with hormones, with life events, with relationship changes, with the accumulated experience of living in a body across decades. Every life stage has its own challenges and, perhaps less discussed, its own gifts.

18.1 Adolescence and Early Experience

The most important thing that can happen during this period is not orgasm. It is the development of a relationship with one's own body as a source of information and pleasure rather than evaluation. Solo exploration, body knowledge, and an early grounding in consent build the foundation for everything that comes later. For young women reading this book: your body is not a problem to be solved. It is an instrument to be learned.

18.2 Young Adulthood (20 to 35)

For many women, this is the period of highest spontaneous desire, most active sexual exploration, and greatest accumulation of partnered experience. It is also frequently the period of most faking, most silence, and most accommodation of a partner's pleasure at the expense of one's own. The investment here is communication and self-knowledge.

18.3 The Parenting Phase

There is perhaps no phase more consistently challenging for female sexuality than the parenting phase. Fatigue is a profoundly effective arousal inhibitor. Body image shifts. Identity reorganises entirely. Spontaneous desire often retreats further. The responsive desire model becomes not just theoretically relevant but practically necessary. Protecting intimate time is one of the most important investments available during this phase.

18.4 Perimenopause and Menopause (40 to 60)

Oestrogen fluctuation produces real physical changes: reduced vaginal lubrication, some tissue thinning, potential changes in arousal timing. These are real and addressable. Lubricants are not a consolation prize. They are a tool that the body has asked for. Pelvic floor work becomes more, not less, important. Hormone replacement therapy is a personal and medical decision that deserves a genuine conversation with a healthcare provider current on the evidence.

> ★ **EVIDENCE NOTE — HRT AND MENOPAUSE: EVOLVING EVIDENCE**
>
> The evidence landscape around hormone replacement therapy has shifted significantly since the early 2000s when the WHI study produced initial concerns (subsequently reanalysed). Current consensus from bodies including the Menopause Society and ISSWSH supports that for many women, particularly those under 60 or within 10 years of menopause onset, HRT offers more benefit than risk. This is a rapidly evolving area. Discuss current guidelines with a menopause-informed healthcare provider.

18.5 Post-Menopause (60 and Beyond)

Many women report the best sex of their lives in their 60s, 70s, and beyond. Pregnancy is no longer a concern. The performance anxiety and body image concerns of younger decades have, for many women, substantially reduced. Decades of accumulated self-knowledge and confidence make up for whatever hormonal shifts have occurred. Sexuality in older women is consistently and poorly represented in culture. It exists. It is significant. It deserves better.

✦ CHAPTER SUMMARY — PLEASURE ACROSS THE LIFESPAN

• Every life stage brings both challenges and gifts to sexual experience
• Adolescence: build body relationship, not orgasm performance
• Young adulthood: invest in communication and self-knowledge — compounding returns
• Parenting phase: protect intimate time; responsive desire becomes essential, not theoretical
• Perimenopause/menopause: changes are real and addressable; discuss HRT with current evidence
• Post-menopause: many women report their best sexual years — representation is failing this demographic

CHAPTER 19

Queer Pleasures & Expanded Relationship Structures

❧

The orgasm gap conversation belongs to every woman. This chapter is specifically for women whose experiences have been largely absent from the mainstream sexuality bookshelf: lesbian and bisexual women, queer women, women in non-monogamous relationships, asexual-spectrum women, and women in relationships that do not conform to the partnered heterosexual script this book has, by necessity, often used as its default frame.

★ EVIDENCE NOTE — LESBIAN ORGASM RATES: STRONG EVIDENCE — AND THE ARGUMENT THIS WHOLE BOOK IS BUILT ON

The 86% orgasm rate among lesbian women (Frederick et al., 2018, Archives of Sexual Behavior) is among the most powerful data points in this entire book. It confirms that the orgasm gap is behavioural and educational, not biological. Lesbian sexual practice tends to prioritise external clitoral stimulation, extended arousal time, and mutual attention — exactly the principles this book advocates.

19.1 Lesbian and Bisexual Women's Sexual Experience

For women who have sex primarily or exclusively with women, the orgasm gap often does not apply in the same way. The elimination of the penetration-as-default script and the presence of a partner who

understands female anatomy through lived experience produces dramatically better outcomes for orgasm frequency. This is not coincidence. It is curriculum.

Bisexual women, however, report some of the most complex sexual experiences in survey data: lower orgasm rates than both lesbian and heterosexual women in some studies, higher rates of sexual anxiety, and a persistent sense of being invisible in sexuality education that is framed for either heterosexual or lesbian women but rarely for both. This book sees you.

19.2 Non-Monogamy and Relationship Diversity

Consensual non-monogamy, including polyamory, open relationships, swinging, and relationship anarchy, involves sexual and romantic structures that challenge the paired-coupling assumption of most sexuality writing. Women in these structures often report higher communication skills around sexuality, given that consent and desire must be negotiated explicitly across multiple partners. The communication frameworks in Chapter 11 apply, and apply with additional nuance: multiple partners mean multiple conversations, multiple sets of preferences, and the need for particularly clear personal self-knowledge.

Jealousy, time negotiation, and emotional complexity are realities of non-monogamous structures that deserve acknowledgment without judgment. So does the finding in some research that women in ethically non-monogamous relationships report higher sexual satisfaction than their monogamous counterparts. This is not an advertisement for non-monogamy. It is an observation that structures

which prioritise explicit communication and mutual desire tend to produce better sexual outcomes.

19.3 Asexual Spectrum Experiences

Asexuality is a sexual orientation characterised by little or no sexual attraction to others. It is not a disorder, not a trauma response, not a phase, and not celibacy by choice. The asexual spectrum includes asexual individuals who experience romantic attraction, demisexual individuals who experience sexual attraction only after deep emotional connection, and graysexual individuals who experience sexual attraction rarely or under specific circumstances.

For asexual-spectrum women who do engage in sexual activity with partners, the frameworks in this book for communication, emotional safety, and consent architecture remain fully relevant. For those who do not, the body-knowledge and self-understanding chapters apply on their own terms. Your experience of your own body is valid regardless of whether you share it.

19.4 Trans and Non-Binary Women's Experiences

Trans women and non-binary people with vulvas are women for the purposes of this book. The anatomy chapters apply as written. For trans women whose anatomy differs from the cisgender female body described in Chapter 2, much of the psychology, communication, and emotional safety content remains directly applicable, with the understanding that individual anatomy varies and that professional support from a trans-affirming sex therapist or healthcare provider is

particularly valuable for navigating the intersection of gender and sexuality.

Gender dysphoria can intersect with sexual experience in complex ways that no generalised book can fully address. What this book can offer is the unambiguous statement that trans women's sexuality and pleasure deserve the same thoughtful, evidence-based attention as anyone else's.

💬 REAL WOMEN, REAL STORIES — CASS, 33, AUSTRALIA

"The thing that nobody ever said to me explicitly until I was well into my thirties was: the way sex works for you is not less real just because it doesn't look like the default story. I'm bisexual, I've been in a polyamorous relationship for six years, and I spent a long time feeling like I was some kind of footnote. I wasn't a footnote. I was just underrepresented. There's a difference."

On being seen in a conversation that usually doesn't include you. Names and identifying details have been changed to protect privacy.

✦ CHAPTER SUMMARY — QUEER PLEASURES & EXPANDED RELATIONSHIP STRUCTURES

• The 86% lesbian orgasm rate is the strongest argument in this book: the gap is educational, not biological
• Bisexual women are underserved by sexuality education that addresses only lesbian or heterosexual experience
• Non-monogamous structures that prioritise communication tend to produce better sexual outcomes
• Asexuality is an orientation, not a disorder, a phase, or a choice
• Trans women's sexuality deserves the same evidence-based attention and respect

CHAPTER 20

Closing the Gap Together

We have covered a lot of ground. Anatomy, techniques, psychology, communication, desire, trauma, disability, queer experience, the full lifespan, the full geography of pleasure. If you have read this far, you now have more comprehensive knowledge of female sexual response than most healthcare providers, most sex educators, and most partners you will ever encounter.

This is not an accident. The information has always existed. What has been lacking is the will to gather it, organise it, and put it in the hands of the people who need it most.

20.1 The Five Commitments

The Knowledge Commitment

I commit to understanding my own anatomy and, if I have a partner, their anatomy too. Knowledge is the first act of care.

The Communication Commitment

I commit to speaking about pleasure before, during, and after sexual encounters as a normal, caring, and welcome act.

The Technique Commitment

I commit to expanding my sexual repertoire beyond default scripts and experimenting with what the evidence actually suggests works.

The Patience Commitment

I commit to understanding that female orgasm is not a performance metric with a success threshold and a time limit. An encounter without orgasm is not a failed encounter.

The Equity Commitment

I commit to treating my partner's pleasure as equal in importance to my own. For partners who have historically prioritised their own pleasure, this is an invitation to expansion. For women who have historically subordinated their pleasure, it is an invitation home.

20.2 A Letter to Partners

If you are reading this book as a partner of someone with a vulva, we want to say something directly to you.

The orgasm gap is not the result of individual failure. It is the result of collective ignorance. No one in this gap is the villain. But you are reading this book. Which means you are not prepared to leave things as they were. That matters more than you might think. The most consistent finding in research on sexual satisfaction is that partners who are curious, attentive, and communicative produce dramatically better outcomes than partners who are technically skilled but emotionally closed. You do not have to have done it right up until now. You just have to be willing to do it differently from here.

20.3 The Gap Closes in Practice

Knowledge is necessary but not sufficient. The gap closes in real encounters, in real relationships, in real bodies, through practice. Through the awkward first conversation about what feels good.

Through the slightly self-conscious reaching for one's own hand during sex. Through the willingness to say that was amazing and I would love more of it.

It closes through anatomy honestly taught. Through techniques that respect the actual structure of female pleasure. Through communication that builds the shared language of desire. Through emotional safety that creates the conditions for genuine surrender. Through the patient, curious, evidence-based understanding that female orgasm is not a mystery. It was never a mystery. It was simply an absence of information.

You now have the information. The rest is practice.
Welcome to the other side of the gap.

APPENDIX A

Master Do's & Don'ts Reference

Universal Do's

• Prioritise clitoral stimulation in every partnered encounter.
• Communicate about pleasure before, during, and after sex.
• Use lubrication generously.
• Allow at least 15 to 20 minutes before attempting penetration.
• Ask for and give specific, real-time feedback.
• Experiment with positions, angles, and stimulation types.
• Treat orgasm as one of many pleasurable outcomes, not the sole measure of success.
• Support solo practice.
• Create emotional safety as a precondition for great sex.
• Discuss desires outside of the bedroom.
• For trauma survivors: communicate safety signals and the right to stop without explanation.
• For partners: presence outperforms expertise. Be here.

Universal Don'ts

• Assume penetration alone will produce orgasm.
• Interpret silence as satisfaction.
• Treat lack of lubrication as lack of interest.
• Rush to penetration before full arousal.
• Change technique constantly. Consistency enables orgasm.
• Make orgasm the sole measure of sexual success.
• Feel competitive with vibrators or other tools.
• Wait for the heat of the moment for difficult conversations.
• Expect physical technique to compensate for emotional distance.
• Fake orgasms as a long-term strategy.
• Push through pain. Pain is information, not a challenge.
• Dismiss pelvic pain conditions. They are medical, not psychological.

APPENDIX B

Technique Combination Strategy Cards

Angling + Pairing

Best in missionary with pillow or modified doggy. The most reliable penetrative-orgasm combination. Start here.

Rocking + Pairing

Best in cowgirl or spooning. Receiver controls both the rocking rhythm and clitoral stimulation simultaneously.

Shallowing + Oral

Receiver on back. Excellent extended foreplay. Delay penetration for maximum effect.

G-Spot + Clitoral (Blended)

Receiver on back, partner using manual stimulation. Fingers to anterior wall plus tongue or thumb on clitoris simultaneously.

Edging + Pairing

Any comfortable position. Approach orgasm via pairing, reduce stimulation three times before allowing release.

Pelvic Floor + Angling

Cowgirl or CAT missionary. Receiver pulses pelvic floor muscles while maintaining angled anterior-wall contact.

Breathwork + A-Spot

Missionary or receiver on back. Deep, slow, conscious breathing during deep penetration at full arousal.

APPENDIX C

Evidence Calibration Guide: What the Research Actually Proves

★ **EVIDENCE NOTE — A NOTE ON DATA CURRENCY**

Sexual behaviour research evolves. Survey statistics on orgasm frequency, prevalence rates, and sexual practice patterns should be understood as current best estimates rather than permanent fixed figures. The directional findings — that a gap exists, that it is educational rather than biological, that communication matters, that clitoral stimulation is primary — are robust across decades of research. Specific percentages may shift as studies are replicated, sample diversity improves, and cultural contexts change. Readers are encouraged to check the research appendix and primary sources for the most recent data.

Well-Established (Strong, Replicated Evidence)

- The orgasm gap exists: women orgasm significantly less frequently than men in heterosexual partnered sex (Frederick et al., 2018; OMGYES 2016, 2020; Herbenick et al., 2010).
- The majority of women require clitoral stimulation to orgasm, and the minority reliably orgasm through penetration alone (Herbenick et al., 2010; Lloyd, 2005).
- The clitoris is a multi-part organ extending internally; complete anatomy accurately mapped in 1998 (O'Connell et al., Journal of Urology).
- Lesbian women have orgasm rates comparable to heterosexual men, strongly indicating the educational/behavioural rather than biological nature of the gap (Frederick et al., 2018).
- Communication with partners is positively and consistently correlated with female orgasm frequency (Mark et al., 2014).
- Trauma affects sexual response through neurological mechanisms, not weakness of character (van der Kolk, 2014).

• SSRI use affects sexual function in 30–70% of users (Clayton et al., 2014).

Moderately Supported (Consistent but Smaller or Fewer Studies)

• The Dual Control Model of sexual response (SES/SIS): well-theorised with strong clinical application; large population-level replications ongoing (Bancroft & Janssen, 2000).
• The A-spot: documented in Chua (1997) with compelling results; less independently replicated than G-spot research.
• Pelvic floor strengthening improving orgasm intensity: supported by physiotherapy literature; not yet the subject of large RCTs.
• Breathwork affecting arousal: supported by nervous system physiology; direct sexual application less studied.
• CAT technique improving orgasm frequency: Pierce et al. (2000); small sample, anatomically sound rationale.

Emerging or Contested (Calibrate Expectations)

• Female ejaculation composition: studies vary; the balance of prostate-fluid and urinary components is not yet settled (Salama et al., 2015). Both positions have published research.
• G-spot as distinct anatomical structure vs. internal clitoris: O'Connell (1998) and Buisson (2021) support the latter; others maintain functional distinctiveness. Practical techniques are the same either way.
• Cervical orgasm: documented through case study and qualitative research; no large controlled studies to date (Komisaruk & Whipple, 2011).
• Exercise-induced orgasm prevalence: documented (Herbenick & Fortenberry, 2011); mechanisms incompletely understood.
• Full-body orgasm via breathwork: supported by clinical observation and tantric traditions; limited formal neuroimaging research.

APPENDIX D

Recommended Resources & Research

Key Research Sources

• OMGYES Research (2016, 2020): omgyes.com. The largest survey study of female pleasure techniques ever conducted.
• Nagoski, Emily. Come As You Are (2015). The foundational accessible text on female sexual response and the Dual Control Model.
• O'Connell, H.E. et al. (1998). Anatomical relationship of the urethra and clitoris. Journal of Urology, 159(6), 1892–1897.
• Frederick, D.A. et al. (2018). Differences in Orgasm Frequency Among Gay, Lesbian, Bisexual, and Heterosexual Men and Women. Archives of Sexual Behavior, 47(1), 273–288.
• van der Kolk, Bessel. The Body Keeps the Score (2014). Essential for understanding trauma and somatic experience.
• Buisson, O. et al. (2021). Ultrasound visualisation of internal clitoral anatomy during arousal. Clinical Anatomy.
• Basson, R. (2002). A model of women's sexual arousal. Journal of Sex & Marital Therapy, 28(1), 1–10.
• Mark, K.P. et al. (2014). Sexual communication in romantic relationships. Archives of Sexual Behavior.

Recommended Books

• Come As You Are by Emily Nagoski
• She Comes First by Ian Kerner
• The Vagina Bible by Jen Gunter
• Becoming Cliterate by Laurie Mintz
• Mating in Captivity by Esther Perel
• The Body Keeps the Score by Bessel van der Kolk
• Women's Anatomy of Arousal by Sheri Winston

• Healing Sex: A Mind-Body Approach to Healing Sexual Trauma by Staci Haines

Global Support Organisations

• OMGYES.com — Evidence-based, touch-interactive video platform for female pleasure techniques.
• AASECT (aasect.org) — Use the therapist finder to locate a sex therapist near you.
• RAINN (rainn.org) — Sexual violence support and resources (US-based with international links).
• ISSWSH (isswsh.org) — International Society for the Study of Women's Sexual Health.
• One Billion Rising (onebillionrising.org) — Global anti-violence network with country-specific resource directories.
• Pelvic Health Solutions (pelvichealthsolutions.ca) — Pelvic floor physiotherapy information and international provider directory.
• ISTSS (istss.org) — International trauma therapist directory.
• Relate (relate.org.uk) — UK-based relationship and sex therapy.
• Family Planning Alliance Australia (fpaa.org.au) — Sexual and reproductive health resources for Australia and the Pacific.

APPENDIX E

Glossary of Key Terms

The following terms are used throughout this book. Where multiple definitions exist in the literature, we use the definition most consistent with current clinical consensus.

Arousal non-concordance

The state in which physical arousal (lubrication, engorgement) and psychological arousal (desire, interest) occur independently of each other.

Clitoral complex (CUV)

The clitorourethrovaginal complex: the full internal clitoral structure including glans, body, crura, and vestibular bulbs, together with the urethra and anterior vaginal wall.

Coital Alignment Technique (CAT)

A missionary position variation in which the penetrating partner positions their pubic bone over the receiver's clitoris and uses rocking rather than thrusting motion.

Dual Control Model

Bancroft and Janssen's theoretical framework describing sexual response as the net result of a sexual excitation system (accelerator) minus a sexual inhibition system (brakes).

Orgasm gap

The well-documented disparity in orgasm frequency between heterosexual women and men during partnered sex, approximately 30 percentage points.

Pairing

The simultaneous addition of clitoral stimulation during penetrative sex.

Pelvic floor

The group of muscles and connective tissue forming the base of the pelvic cavity, which contract rhythmically during orgasm.

Responsive desire

A desire pattern in which sexual interest emerges in response to erotic stimulation rather than appearing spontaneously before stimulation.

Sensate focus

A structured touch exercise programme developed by Masters and Johnson, used in sex therapy to reduce performance anxiety and build body awareness.

Spectatoring

The habit of mentally stepping outside one's own body during sex to evaluate oneself from an imagined external perspective.

Tenting effect

The lifting and expansion of the vaginal canal and uterus during full arousal.

Vasocongestion

The engorgement of genital tissues with blood during arousal, affecting the clitoris, vestibular bulbs, and vaginal walls.

Vaginismus

Involuntary spasm of the pelvic floor muscles that makes penetration painful or impossible; a medical condition treatable with pelvic floor physiotherapy.

Vulvodynia

Chronic vulval pain without an identifiable cause; a medical condition with evidence-based treatments available.

APPENDIX F

Notes & References

The following is a consolidated reference list for key claims made throughout this book. Readers are encouraged to seek the full texts for deeper engagement with the primary research.

Chapter 1 — The Orgasm Gap

• Frederick, D.A. et al. (2018). Differences in Orgasm Frequency Among Gay, Lesbian, Bisexual, and Heterosexual Men and Women. Archives of Sexual Behavior, 47(1), 273–288.
• Herbenick, D. et al. (2010). Sexual Behavior in the United States: Results from a National Probability Sample. Journal of Sexual Medicine, 7(s5), 255–265.
• Lloyd, E.A. (2005). The Case of the Female Orgasm. Harvard University Press.
• Muehlenhard, C.L. & Shippee, S.K. (2010). Men's and Women's Reports of Pretending Orgasm. Journal of Sex Research, 47(6), 552–567.
• OMGYES Research (2016, 2020). omgyes.com. Methodology available on site.

Chapter 2 — Anatomy

• Chivers, M.L. et al. (2010). Agreement of Self-Reported and Genital Measures of Sexual Arousal. Archives of Sexual Behavior, 39(1), 5–56.
• O'Connell, H.E. et al. (1998). Anatomical Relationship of the Urethra and Clitoris. Journal of Urology, 159(6), 1892–1897.

Chapters 4–7 — Techniques

• Buisson, O. et al. (2021). The Nature of Female Orgasm as Revealed by Ultrasound Visualisation During Coition. Clinical Anatomy.
• Pierce, A.P. et al. (2000). Coital Alignment Technique and Directed Masturbation. Journal of Sex & Marital Therapy, 26(2), 117–131.

Chapters 8–11 — Psychology & Communication

• Bancroft, J. & Janssen, E. (2000). The Dual Control Model of Male Sexual Response. Neuroscience and Biobehavioral Reviews, 24(5), 571–579.
• Basson, R. (2002). A Model of Women's Sexual Arousal. Journal of Sex & Marital Therapy, 28(1), 1–10.
• Mark, K.P. et al. (2014). Sexual Communication, Sexual Satisfaction, and Relationship Satisfaction. Archives of Sexual Behavior.
• Vannier, S.A. & O'Sullivan, L.F. (2012). Sex Without Desire. Journal of Sex & Marital Therapy, 37(3), 179–197.

Chapters 12–14 — Erogenous Map

• Chua, C.A. (1997). A Proposed Model for the Activation of Sensory Nerve Endings in the Glans Clitoridis. Medical Hypotheses, 48(3), 214–216.
• Komisaruk, B.R. & Whipple, B. (2011). Non-Genital Orgasms. Sexual and Relationship Therapy, 26(4), 356–372.

Chapter 15 — Trauma

• van der Kolk, B. (2014). The Body Keeps the Score: Brain, Mind, and Body in the Healing of Trauma. Viking.

Chapter 16 — Disability

• Bergeron, S. et al. (2015). The Treatment of Vulvodynia: A Review. Journal of Lower Genital Tract Disease.
• Rosenbaum, T.Y. & Owens, A. (2008). The Role of Pelvic Floor Physical Therapy in the Treatment of Pelvic and Genital Pain-Related Sexual Dysfunction. Journal of Sexual Medicine, 5(3), 513–523.

Chapter 17 — Partners

• Kleinplatz, P.J. et al. (2009). The Components of Optimal Sexuality. Sexual and Relationship Therapy, 24(1), 3–13.

Chapter 18 — Lifespan

• Waldinger, M.D. et al. (2005). A Multinational Population Survey of Intravaginal Ejaculation Latency Time. Journal of Sexual Medicine, 2(4), 492–497.

❧

THE ORGASM GAP

First Edition

"The gap closes here."

❧